

Patching up the monopole potential

Ashok Chatterjee* and P. Mitra†
 Saha Institute of Nuclear Physics
 Block AF, Bidhannagar
 Calcutta 700 064, INDIA

hep-th/0312272

Abstract

It is well known that a vector potential cannot be defined over the whole surface of a sphere around a magnetic monopole. A recent claim to the contrary is shown to have problems. It is explained however that a potential of the proposed type works if two patches are used instead of one. A general derivation of the Dirac quantization condition attempted with a single patch is corrected by introducing two patches. Further, the case of more than two patches using the original Wu-Yang type of potential is discussed in brief.

While magnetic monopoles have not been seen experimentally, they have continued to be of interest to theoreticians. In standard electrodynamics, magnetic monopoles are not admitted, and the magnetic field has zero divergence. If a magnetic monopole is present at a point, the divergence is a delta function with support at that point, and it is not possible to introduce a vector potential which is nonsingular everywhere. Dirac[1] found that there is at least a string of singularities from that point to infinity, though the location of the string is arbitrary, like a branch cut. Instead of working with a string, Wu and Yang [2] showed that it may be more convenient to work with two different nonsingular potentials on two overlapping patches, with a gauge transformation connecting them in the overlap. In both approaches the quantum theoretic description of an electrically charged particle requires the quantization of its charge, if a monopole is assumed to exist.

Recently there has been a claim [3] about the construction of a nonsingular vector potential for the monopole field using a *single* patch. This could only be possible if the basis of the Wu-Yang formalism were incorrect, and indeed that is what has been asserted. One of our aims is to reexamine this question and to show that the *single*-patch potential does not work. The use of two patches,

*e-mail ashok@theory.saha.ernet.in

†e-mail mitra@theory.saha.ernet.in

with a potential in the ϕ direction [2], is standard. We show how the vector potential of [3], which is in the θ direction, can be extended to two patches, where it makes sense. A general argument for the Dirac quantization, presented for a single patch in [3], also does not work. We show how it can be corrected for two patches. We then discuss the possibility of using the ϕ -directional potential of [2] in *more than two* patches: although two patches are enough, one can use more, contrary to claims in [3]. There arise interesting issues of consistency in this situation, but the standard quantization condition continues to hold.

The potential proposed [3] for a monopole of strength g at the origin is locally given by the 1-form

$$A = -gf(\phi) \sin \theta d\theta. \quad (1)$$

This leads to the 2-form field

$$F = dA = \frac{g}{r^2} f'(\phi) r d\theta \wedge r \sin \theta d\phi, \quad (2)$$

which reduces to the standard expression for the magnetic field for the monopole at the origin if and only if f' is unity, *i.e.*,

$$f(\phi) = \phi + \text{constant}. \quad (3)$$

This choice makes f multiple valued in the sense that ϕ and $\phi + 2\pi$ give different values. If the angle were taken mod 2π , the derivative would no longer be unity: the mod involves a sawtooth function, whose derivative is a sum of delta functions with supports at integral values of $\phi/2\pi$, so that the magnetic field would have singularities there. If ϕ is restricted to an interval of size 2π , the potential becomes discontinuous at the boundaries of the interval. In general, for any f , one may note that the line integral on a closed curve C with $\theta = \text{constant}$ on the sphere $r = \text{constant}$ is well defined and given by

$$\oint_C A = 0. \quad (4)$$

If Stokes' theorem, which is valid for a nonsingular potential, is applicable, it implies that the flux through a surface bounded by C is zero. This is obviously not the case for the magnetic field due to a monopole. Therefore, the potential cannot be nonsingular. It should be pointed out that the line integral has been proposed [3] to be replaced by an integral over a complicated path so that it ceases to be zero, but such a replacement can be justified only if the potential is defined on two patches – as discussed below – instead of one. Otherwise it is *ad hoc* to replace the well-defined line integral by something which agrees with the answer expected from Stokes' theorem.

In fact, the impossibility of constructing a vector potential using a single patch is a theorem in [2]. The idea is that the flux through the part of the sphere $r = \text{constant}$ with $\theta < \theta_0$, where $0 < \theta_0 < \pi$, increases as θ_0 increases from zero to π , reaching the total flux $4\pi g$ in the limit. However, for each θ_0 , the flux is also equal to $\oint_{C[\theta_0]} A$, where $C[\theta_0]$ is the curve $\theta = \theta_0$ on the sphere, if

A can be defined over the whole sphere. In the limit $\theta_0 \rightarrow \pi$, this curve reduces to a point, so that the line integral of a nonsingular A must vanish, yielding a contradiction $4\pi g \stackrel{?}{=} 0$ for $g \neq 0$.

This contradiction was dismissed in [3], on the ground that the curve $C[\theta_0]$ is also the boundary of another part of the spherical surface, namely that with $\theta > \theta_0$, which vanishes in the limit $\theta_0 \rightarrow \pi$, so that the flux through it is zero. But this is a red herring. For a potential which is regular everywhere, the line integral on the curve must be equal to the surface integral on *each* of the two parts (taken in the appropriate sense). The contradiction between the flux on the whole sphere and the line integral on a curve of vanishing length can be recast as a mismatch between the fluxes on the whole surface and its null complement. This contradiction cannot be removed without removing the monopole if one persists in the attempt to use a single patch. One needs at least two patches to construct a nonsingular A on a sphere $r = \text{constant}$.

As is well known, one possibility [2] is to take the two patches as $\theta < \theta_1$ and $\theta > \theta_2$ with $0 < \theta_2 < \theta_1 < \pi$. If the line integral on a curve with constant θ and varying ϕ is to be nonzero, the potential 1-form must have a $d\phi$ piece. Equating the line integral to the surface integral in the first patch, one finds this piece to be

$$A^{(1)} = g(1 - \cos \theta)d\phi \quad 0 < \theta < \theta_1. \quad (5)$$

Similarly the surface integral appropriate for the second patch yields

$$A^{(2)} = g(-1 - \cos \theta)d\phi \quad \theta_2 < \theta < \pi. \quad (6)$$

In the overlapping region, where both of these expressions are valid, they are clearly unequal, showing the impossibility of having a single patch, but their difference is closed, and as expected from Poincaré's lemma, locally exact:

$$A^{(1)} - A^{(2)} = d\chi, \quad (7)$$

with

$$\chi = 2g\phi. \quad (8)$$

Although χ is multiple-valued, $d\chi$ is well defined, so there is no problem in the context of the classical theory. However, if there is a particle with electric charge e in this magnetic field, and it is described by a quantum mechanical wavefunction, the gauge transformation of this wavefunction involves the factor $\exp(ie\chi/\hbar c)$ and this has to be single-valued. This requires

$$2ge = n\hbar c \quad n \text{ integral}, \quad (9)$$

which is the Dirac quantization condition. We have reviewed this well known analysis in some detail because we want to extend it in two ways: to the potential proposed in [3] and to the case of more than two patches which has also been questioned.

If we seek to use the kind of potential proposed in [3], we have to take patches in the ϕ direction rather than the θ direction. Thus, using two patches as before, we may set

$$A^{(1)} = -g\phi \sin \theta d\theta \quad \phi_1 < \phi < \phi'_1, \quad (10)$$

and

$$A^{(2)} = -g\phi \sin \theta d\theta \quad \phi_2 < \phi < \phi'_2, \quad (11)$$

with

$$\phi_2 < 0 < \phi_1 < \phi'_2 < \phi_2 + 2\pi < \phi'_1 < 2\pi. \quad (12)$$

The two patches together cover the full region $(0, 2\pi)$ with overlaps $R_1 \equiv (\phi_1, \phi'_2)$ and $R_2 \equiv (\phi_2 + 2\pi, \phi'_1)$. In these overlapping regions, the two potentials are gauge related:

$$A^{(1)} - A^{(2)} = \begin{cases} 0 & \phi \text{ in } R_1 \\ 2\pi g d \cos \theta & \phi \text{ in } R_2 \end{cases}. \quad (13)$$

The 2π comes from the fact in R_2 , the coordinate ϕ has values differing by 2π in the two patches. One may write these in the form (7) with

$$\chi = \begin{cases} 0 & \phi \text{ in } R_1 \\ 2\pi g(\cos \theta + p) & \phi \text{ in } R_2 \end{cases}, \quad (14)$$

where p is a constant. For θ different from $0, \pi$, the regions R_1, R_2 are non-overlapping. However, when θ goes to either of the limiting values, ϕ is undefined, and the two regions cannot be distinguished. Clearly, the two expressions for χ cannot be made to agree at both $\theta = 0$ and $\theta = \pi$ by any choice of p , but by choosing it to be equal to either -1 or +1, they can be made to agree at either $\theta = 0$ or $\theta = \pi$. At the other point, χ has two unequal values, but as in the earlier discussion, the gauge transformation $\exp(i e \chi / \hbar c)$ of the wavefunction of a particle with electric charge e becomes single-valued if and only if the Dirac condition (9) holds. In this way, the θ -directional potential [3] can be made to work with two patches, and the Dirac condition emerges as the consistency condition. The evaluation of a line integral on a loop with constant θ can be carried out now in the manner indicated in [3], the deformation of path suggested there being equivalent to the necessary gauge transitions between the two patches.

An attempt was made in [3] to derive the Dirac condition without invoking any specific form of the potential, but it is based on a single patch, which as we know, is impossible. However, it is not difficult to repair the derivation using two patches, which we shall now do. Suppose the sphere around the monopole is covered by two overlapping patches and nonsingular potentials $A^{(1)}, A^{(2)}$ defined on them to describe the monopole magnetic field. Then in the overlapping region,

$$F = dA^{(1)} = dA^{(2)}. \quad (15)$$

Thus in this region the two potentials must again be related by (7), with a χ determined by $A^{(1)}, A^{(2)}$. Now consider a closed loop C which lies entirely in the overlap region and is non-contractible within the overlap. Then the two surfaces S_1, S_2 bounded by C lie in the two patches and cover the full sphere. The application of Stokes' theorem to the two patches leads to the two separate equations

$$\begin{aligned}\oint_C A^{(1)} &= \int_{S_1} F, \\ \oint_C A^{(2)} &= - \int_{S_2} F,\end{aligned}\tag{16}$$

where the difference in orientation of the two surfaces relative to a given direction of traversal of C has been taken into account. The above two equations, on subtraction, lead to the result that

$$\Delta\chi \equiv \oint_C d\chi = \int_{S_1 \cup S_2} F = 4\pi g \neq 0.\tag{17}$$

This shows that the gauge transformation function χ is not single-valued, but the relevant factor for the wavefunction of a particle of charge e is $\exp(ie\Delta\chi/\hbar c)$ and it is single-valued if and only if the Dirac condition (9) is satisfied. This demonstration does not rely on explicit formulas for A in the ϕ or θ directions. But it does involve A defined on two patches. A gauge invariant derivation without introducing A was given in [4] on the basis of the assumption that translation operators are associative.

Finally we turn to the case with more than two patches. The extension of the analysis of [2] using two patches to the case of more patches has been questioned [3]. A clarification is in order here. If there are N patches (N integral), the first patch has the same potential (5) as before, and in the last patch one has

$$A^{(N)} = g(-1 - \cos\theta)d\phi \quad \theta_N < \theta < \pi.\tag{18}$$

but now there are other patches in between, and for these patches, the potentials are of the form

$$A^{(r)} = g(a_r - \cos\theta)d\phi \quad \theta_r < \theta < \theta_{r'},\tag{19}$$

where the angles θ_r and $\theta_{r'}$ for different r are such that the successive patches overlap. The constants a_r can be fixed only for $r = 1, N$ by requiring regularity at 0 and π for the potentials on the two extreme patches:

$$a_1 = 1, \quad a_N = -1.\tag{20}$$

Classically, once again, there is no problem with gauge equivalence of these expressions in overlapping regions, but if there is an electrically charged particle being treated quantum mechanically, one needs at least the conditions

$$(a_r - a_{r+1})ge = n_r\hbar c \quad n_r \text{ integral}\tag{21}$$

from successive overlapping patches, as in (9). If there are further overlaps, for example between the r th and the s th with $|r - s| \neq 1$, there may apparently be further conditions, but they are not independent and can be obtained from those in (21). In any case, by adding up the conditions indicated above, and using the values of a_1, a_N , one gets the Dirac condition (9) again, with

$$n = (n_1 + \dots + n_{N-1}). \quad (22)$$

While this is a straightforward generalization of the case with two patches, there is something new here: the a_r for $2 \leq r \leq N-1$ are not determined by regularity conditions. They are not totally arbitrary, however. In fact, the allowed values are the rational numbers

$$a_r = 1 - \frac{2(n_1 + \dots + n_{r-1})}{(n_1 + \dots + n_{N-1})}, \quad (23)$$

as follows from the overlap conditions. We add however that the number and allowed values of a_r are dictated by our choice of patches and the physics of the problem is entirely contained in the Dirac condition (9).

To summarize, we have shown that the single-patch potential claimed to describe a magnetic monopole does not work. The standard proof that at least two patches are needed is valid. A nonsingular version of the θ -directional potential using two patches has been worked out. A general derivation of the Dirac condition using two patches has been presented following the earlier single-patch attempt. More than two patches can also be used, and we have worked out the possible forms of the potential in the different patches using the more familiar ϕ -directional potential [2].

Acknowledgment

We would like to thank our colleague Palash Baran Pal for many discussions on [3]. We are also grateful to Amit Ghosh and Parthasarathi Majumdar for several conversations on the subject.

References

- [1] P. A. M. Dirac, Proc. Roy. Soc. (London) **A133** (1931) 60
- [2] T. T. Wu and C. N. Yang, Phys. Rev. **D 12** (1975) 3845
- [3] R. K. Ghosh and P. B. Pal, Phys. Lett. **B 551** (2003) 387
- [4] R. Jackiw, Phys. Rev. Lett. **54** (1985) 159